

## Juniperus scopulorum/Oryzopsis micrantha Woodland

COMMON NAME Rocky Mountain Juniper / Little-seed Mountain Ricegrass Woodland  
SYNONYM Rocky Mountain Juniper / Little-seed Ricegrass Woodland  
PHYSIOGNOMIC CLASS Woodland (II)  
PHYSIOGNOMIC SUBCLASS Evergreen Woodland (II.A)  
PHYSIOGNOMIC GROUP Temperate or subpolar needle-leaved evergreen woodland (II.A.4)  
PHYSIOGNOMIC SUBGROUP Natural/Semi-natural (II.A.4.N)  
FORMATION Round-crowned temperate or subpolar needle-leaved evergreen woodland (II.A.4.N.a)  
ALLIANCE JUNIPERUS SCOPULORUM WOODLAND ALLIANCE

CLASSIFICATION CONFIDENCE LEVEL 1

USFWS WETLAND SYSTEM Terrestrial

### RANGE

#### **Theodore Roosevelt National Park**

Rocky Mountain juniper / little-seed ricegrass woodlands are predominantly found on slopes in both the North and South Units. Such sites occur throughout the scoria hills and in the transition zones of the upland grassland and old river terrace regions. This association is also sometimes found at the upper reaches of north facing hardwood draws in close association with *Fraxinus pennsylvanica*.

#### **Globally**

This community is found in southeastern Montana, southwestern North Dakota, western South Dakota, eastern Wyoming, and western and central Nebraska.

### ENVIRONMENTAL DESCRIPTION

#### **Theodore Roosevelt National Park**

Slope and aspect appear to play major roles in the extent and development of these woodlands. The soils on such steep slopes tend to be poorly developed clay, clay loams, and scoria slopes (Girard 1985). This alliance is frequently used by native ungulates (bison, elk, and deer) for lounging. Well-worn trails are common on many sites. Stumps from historic logging were also quite common in many of these woodlands.

#### **Globally**

This community typically occurs on moderate to steep (16-70%) north-facing slopes, but can occur on a variety of aspects (Johnston 1987, Von Loh *et al.* 1999). The soils are poorly developed, shallow, loamy sands, sandy loams, and clay loams, sometimes with high gravel content. These woodlands are frequently associated with outcrops of sandstone (DeVelice *et al.* 1995) or scoria and clay slopes (Girard *et al.* 1989).

### MOST ABUNDANT SPECIES

#### **Theodore Roosevelt National Park**

<u>Stratum</u>	<u>Species</u>
Tree Canopy	<i>Juniperus scopulorum</i> , <i>Fraxinus pennsylvanica</i>
Short Shrub	<i>Prunus virginiana</i> , <i>Symphoricarpos occidentalis</i> , <i>Rhus trilobata</i> , <i>J. scopulorum</i>
Herbaceous	<i>Oryzopsis micrantha</i> , <i>Maianthemum stellatum</i> , <i>Elymus virginicus</i>

#### **Globally**

<u>Stratum</u>	<u>Species</u>
Tree canopy	<i>Juniperus scopulorum</i> , <i>Juniperus virginiana</i>
Short shrub	<i>Rhus trilobata</i> , <i>Symphoricarpos occidentalis</i>
Forb	<i>Campanula rotundifolia</i> , <i>Galium boreale</i> , <i>Maianthemum stellatum</i>
Graminoid	<i>Oryzopsis micrantha</i>

### CHARACTERISTIC SPECIES

#### **Theodore Roosevelt National Park**

*Juniperus scopulorum*, *Oryzopsis micrantha*

#### **Globally**

*Juniperus scopulorum*, *Juniperus virginiana*, *Oryzopsis micrantha*

**USGS-NPS Vegetation Mapping Program**  
**Theodore Roosevelt National Park**

---

**VEGETATION DESCRIPTION**

***Theodore Roosevelt National Park***

This community is densely wooded with interlocking canopies of *Juniperus scopulorum*. Mean foliar cover of *J. scopulorum* is about 41%. *Fraxinus pennsylvanica* is a common, but low density associate, especially on the upper reaches of upland draws. *Prunus virginiana* is a frequent understory shrub, often occurring in dense patches. *Oryzopsis micrantha* (= *Piptatherum micranthum*) is the major characteristic herbaceous species. *Smilacina stellata* (= *Maianthemum stellatum*) and a moss (probably *Thuidium abietinum*, Girard 1985) are frequent herbaceous associates.

**Globally**

This woodland community is dominated by small *Juniperus scopulorum* trees through most of its range, and is replaced by *J. virginiana* and introgressant hybrids in the eastern portion of its range in Nebraska (Kaul *et al.* 1983). Some stands contain *Fraxinus pennsylvanica*. Most of these trees are 10-20 cm dbh and 4-6 meters tall (Nelson 1961, Hansen *et al.* 1984). Some trees can be up to 30-40 cm dbh. The basal area has been reported at 22-29 m<sup>2</sup>/ha in North Dakota (Hansen *et al.* 1984) and up to 22-41 m<sup>2</sup>/ha in southeastern Montana and northwestern South Dakota (Hansen and Hoffman 1988). Tree canopy is moderate to dense. In North Dakota, Girard *et al.* (1989) measured densities of 975 trees/ha. Where the canopy is dense the shrub and herbaceous strata are poorly developed. Where the canopy is less full, shrubs and herbaceous species are more abundant. On 7 stands in southwest North Dakota mosses and lichens covered 72% of the ground surface, shrubs covered 17.4%; graminoids - 69.1%; forbs - 9.4% (Hansen *et al.* 1984). Three stands in southeastern Montana had less coverage in each strata (Hansen and Hoffman 1988). Among the shrubs that may be found in this community are *Juniperus communis*, *J. horizontalis*, small *J. scopulorum* or *J. virginiana*, *Mahonia repens*, *Pentaphylloides floribunda*, *Prunus virginiana*, *Rhus trilobata*, *Ribes aureum*, *R. cereum*, *Rosa woodsii*, *Symphoricarpos albus*, and *S. occidentalis*. Typical herbaceous species include *Anemone patens*, *Antennaria microphylla*, *Campanula rotundifolia*, *Carex inops* ssp. *heliophila*, *Chenopodium fremontii*, *Elymus lanceolatus*, *E. trachycaulus*, *Galium boreale*, *Geum triflorum*, *Koeleria macrantha*, *Oryzopsis micrantha*, *Maianthemum stellatum*, *Parietaria pensylvanica*, and *Taraxacum officinale*. *Acer negundo* and *Fraxinus pennsylvanica* saplings are sometimes found in depressions where soil and moisture accumulate.

**OTHER NOTEWORTHY SPECIES**

**CONSERVATION RANK** G3. A number of sites have been impacted by cutting for fenceposts or railroad ties. Fire suppression may increase the extent of the community within its range.

**DATABASE CODE** CEGL000747

**SIMILAR ASSOCIATIONS**

**REFERENCES**

- DeVelice, R.L., S.V. Cooper, J.T. McGarvey, J. Lichthardt, and P.S. Bourgeron. 1995. Plant communities of northeastern Montana: A first approximation. Montana Natural Heritage Program, Helena, MT. 116 p.
- Girard, M.M. 1985. Native woodland ecology and habitat type classification of southwestern North Dakota. Ph.D. Dissertation. North Dakota State University, Fargo. 314 pp.
- Girard, M.M., H. Goetz, and A.J. Bjugstad. 1989. Native woodland habitat types of southwestern North Dakota. Research Paper RM-281. USDA Forest Service, Rocky Mountain Forest and Range Experiment Station, Fort Collins, CO. 36 p.
- Godfread, C. 1994. The vegetation of the Little Missouri Badlands of North Dakota. In Proceedings of the Leafy Spurge Strategic Planning Workshop, March 29-30, Dickinson, ND. Pp 17-24.
- Hansen, P.L., G.R. Hoffman, and A.J. Bjugstad. 1984. The vegetation of Theodore Roosevelt National Park, North Dakota: A habitat type classification. U. S. Dep. Agric., For. Serv., Rocky Mt. For. and Range Exp. Sta., Gen. Tech. Rep. RM-113. Fort Collins, CO. 35 p.
- Hansen, P.L. 1985. An ecological study of the vegetation of the Grand River/Cedar River, Sioux, and Ashland Districts of the Custer National Forest. Unpublished dissertation, South Dakota State University. 257 pp.
- Hansen, P.L. and G.R. Hoffman. 1988. The vegetation of the Grand River/Cedar River, Sioux, and Ashland Districts of the Custer National Forest: a habitat type classification. USDA Forest Service General Technical Report RM-157, Rocky Mountain Forest and Range Experiment Station, Fort Collins, CO.
- Johnston, B.C. 1987. Plant associations of region two: potential plant communities of Wyoming, South Dakota, Nebraska, Colorado, and Kansas. Edition 4. USDA Forest Service, Rocky Mountain Region. R2-Ecol-87-2. 429 pp.
- Kaul, R.B., Challaiah and K.H. Keeler. 1983. Effects of grazing and juniper-canopy closure on the prairie flora in Nebraska high-plains canyons. Proceedings of the Seventh North American Prairie Conference (1980):95-105.
- Nelson, J.R. 1961. Woody plant communities in the badlands of western North Dakota. Proc. North Dakota Acad. Sci. 15:42-44.